



# New phenomena searches at Tevatron Run I

G.-S. Muanza

## ► To cite this version:

G.-S. Muanza. New phenomena searches at Tevatron Run I. Rencontres de Moriond: QCD and Hadronic Interactions at High Energy 37, Mar 2002, Les Arcs, France. pp.20. in2p3-00013755


**HAL Id: in2p3-00013755**

**<https://hal.in2p3.fr/in2p3-00013755>**

Submitted on 22 May 2003

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# New Phenomena Searches at TEVATRON Run I



Steve Muanza, IPN Lyon

On behalf of CDF and D0 collaborations

## Outline

- Run I at the TEVATRON
- Search for SUGRA:
  - dilepton channel at CDF
  - electron channel at D0
- Search for large ED:
  - monojet channel at D0
  - diphoton channel at CDF
- Conclusions

# Collider in Run I



Proton-Antiproton  
Collisions

$\sqrt{s}$	$L$	$\Delta t_x$	Period
$1.8 TeV$	$2 \times 10^{31} cm^{-2} s^{-1}$	$3.5 \mu s$	1992-1996


# Detectors in Run I

Muon:

$$\sigma(r\phi) = 250 \mu\text{m} (\text{CMU})$$

## CDF Detector

- Silicon Vertex Detector
- Large Magnetic Tracking Volume
- Non-Magnetic Central Muon Spectrometer (Forward Toroid)



Calorimeter :

$$\frac{\sigma_{EM}}{E} = \sqrt{\left(\frac{13.5\%}{\sqrt{E_T}}\right)^2 + (2\%)^2}$$

$$\frac{\sigma_{HAD}}{E} = \sqrt{\left(\frac{75\%}{\sqrt{E_T}}\right)^2 + (3\%)^2}$$

Tracking :


$$\sigma(r\phi) = 17 \mu\text{m} (\text{SVX})$$

$$\sigma(z) = 1 \text{ mm} (\text{VTX})$$

$$\frac{\Delta p_T}{p_T} \approx (0.1\%) \times p_T (\text{SVX} + \text{CTC})$$

# Detectors in Run I

- Good Calorimetry: hermeticity, resolution
- Iron Toroid Muon Spectrometer, Large Acceptance



**DØ Detector**

Muon :

$$|\eta| < 3.3$$

$$\frac{\Delta p}{p} = 0.2 \oplus 0.003 p$$

Calorimeter :

$$\frac{\sigma_{EM}}{E} = \frac{15\%}{\sqrt{E}} + 0.3\%$$

$$\frac{\sigma_{HAD}}{E} = \frac{45\%}{\sqrt{E}} + 4\%$$

Tracking :

$$\sigma(\text{vertex}) = 6 \text{ mm}$$

$$\sigma(r\phi) = 60 \mu\text{m (VTX)}$$

$$= 180 \mu\text{m (CDC)}$$

$$\sigma(r\phi) = 200 \mu\text{m (FDC)}$$

# SUGRA Models

## RPC Gravity Mediated SUSY Breaking (MSSM, mSUGRA)

### MSSM Particle Content:

S=1	S=1/2	S=0
	$l$	$\tilde{l}$
	$q$	$\tilde{q}$
$g$	$\tilde{g}$	
$W^\pm$	$\tilde{\chi}_1^\pm$ $\tilde{\chi}_2^\pm$	$\phi_1^+$ $\phi_2^-$
$\gamma$	$\tilde{\chi}_3^0$ $\tilde{\chi}_4^0$	$\phi_1^0$ $\phi_2^0$
$Z$	$\underline{\tilde{\chi}_1^0}$ $\tilde{\chi}_2^0$	

$$M_{\tilde{t}}^2 = \begin{pmatrix} m_{\tilde{t}_L}^2 + m_t^2 & m_t (A_t - \mu/\tan\beta) \\ m_t (A_t - \mu/\tan\beta) & m_{\tilde{t}_R}^2 + m_t^2 \end{pmatrix}$$

### MSSM Phenomenology:

Parameters:

- $m_{1/2}$  : universal gaugino mass
- $A_0$  : universal trilinear coupling
- $\mu$  : higgsino mass parameter
- $\tan\beta$  : ratio of 2 Higgs doublets vev
- $m_0$  : universal scalar mass

R Parity:  $R_P = (-1)^{L+2S+3B}$

Conservation:

- Sparticles pair production
- LSP  $\tilde{\chi}_1^0$  stable  $\Rightarrow E_T^{miss}$  signature


# SUGRA in dileptons at CDF

## Gluino Pair Production

LO contributions:

$$gg \rightarrow \tilde{g}\tilde{g}$$

$$q\bar{q} \rightarrow \tilde{g}\tilde{g}$$



## Gluino and Stop Decays

$$\tilde{g} \rightarrow t\bar{t}_1 / \bar{t} t_1$$

$$\tilde{g} \rightarrow t\bar{t}_1 / \bar{t} t_1$$

- gluino: Majorana sparticle

$$\tilde{t}_1 \rightarrow c \tilde{\chi}_1^0 \quad (\text{stop NLSP})$$

$$t \rightarrow W^\pm b \rightarrow l^\pm \nu b$$

## Final State Topology

Top dilepton events

Like-sign:  $l^\pm l^\pm + jets + E_T^{miss}$

(Opp-sign:  $l^\mp l^\pm + jets + E_T^{miss}$ )

## Main SM Backgrounds

- **Very low**
- Fake lepton ( $W^\pm + \geq 3 \text{ jets}$ )
- SM Top pair production
- Diboson production


# SUGRA in dileptons at CDF

## Analysis

$$\int L dt = 106.1 \text{ pb}^{-1}$$

- **Trigger:** L3 inclusive electron or muon (isolation,  $p_T > 13\text{-}15 \text{ GeV}$ ,...)
- **Lepton ID:** track matching (e), shower shape (e), track-calor. isol. (both),
- **Remove:** conversions (e), cosmics (mu), Z peak (both), radiative Z
- **E<sub>T</sub>miss:**  $E_T^{\text{miss}} > 25 \text{ GeV}$  (isolated)
- **Jets:** at least 2 central jets with  $p_T > 10 \text{ GeV}$

## Selection Results





# SUGRA in dileptons at CDF

## Interpretation of Run I Result

Preliminary 95% *CL Limits*

- Limit on cross section:

$$\sigma_{\tilde{g}\tilde{g}} < 15.1 - 18.0 \text{ pb}$$
$$(200 \leq m_{\tilde{g}} \leq 320 \text{ GeV})$$

$$(BR(\tilde{g} \rightarrow t \tilde{t}) = 1)$$

- Mass limit:


$$m_{\tilde{g}} > 180 \text{ GeV}$$

# mSUGRA in electrons at D0

## Gluino Pair Production/Decay

$$gg \rightarrow \tilde{g}\tilde{g}$$

$$q\bar{q} \rightarrow \tilde{g}\tilde{g}$$



## Final State Topology

$$e^{\pm} + (\geq 4 \text{ jets}) + E_T^{\text{miss}}$$

## Motivation

- Signal sensitive to high and moderate  $m_0$  region: complement to (dilepton and jets)+ $E_T^{\text{miss}}$

## Main SM Backgrounds

$$W^{\pm} + \text{jets}, t\bar{t}, QCD$$

# mSUGRA in electrons at D0

Selection: NN (9-18-1)


$$\int L dt = 94 \text{ pb}^{-1}$$

Preselection

$$E_T(e^\pm) > 20 \text{ GeV}$$


$$E_T(\text{jets}) > 15 \text{ GeV}$$

$$E_T^{\text{miss}} > 25 \text{ GeV}$$




# mSUGRA in electrons at D0

## Results



- $t\bar{t}$  :  $16.8 \pm 5.2$
- $WW + \geq 2$  jets :  $1.4 \pm 0.3$
- QCD Multijet:  $19.1 \pm 4.7$
- $W + \geq 4$  jets :  $43.0 \pm 7.6$
- **Expected Events:  $80.3 \pm 10.4$**
- **Observed Events: 72**

## mSUGRA Limit



# Large Extra Dimensions

Theoretical Foundation:

- Quantum gravity scale  $M_D$  can be  $O(1 \text{ TeV})$ , solution to hierarchy problem


“Modified” Gravity:

$$\frac{4-D}{G_N^{-1} \sim M_{Pl}^2} \longrightarrow \frac{(4+n)-D}{G_N^{-1} \sim R^n M_D^{2+n}}$$

R: compactification (torus) radius

Phenomenology:

- SM particles (open strings): confined to (3+1)-D brane
- Graviton (closed strings): can propagate in the bulk (“weakness of gravity”)
- Equivalence: massive gravitons (KK) in 4-D and massless gravitons (4+n)-D
- Ref: Giudice, Rattazzi, Wells, NPB 544 (1999)
- Ref: Arkani-Hamed, Dimopoulos, Dvali, PLB 429 (1998)
- Ref: Antoniadis, PLB 246 (1990)



# Large ED Searches at Colliders

## ED Nbers and Radii

$$M_D \approx 1 \text{ TeV}$$


$$R = \begin{cases} 1.2 \times 10^{12} m & n = 1 \\ 0.48 \text{ mm} & n = 2 \\ 3.6 \text{ nm} & n = 3 \\ 9.7 \times 10^{-12} m & n = 4 \end{cases}$$

Eöt-Wash Experiment  
 Test Newton's Law down to  
 $O(200 \mu m)$   
 Ref: Adelberger et al, PRL 86 (2001)

## Production Mechanisms


### Real Gravitons

Monojets at hadron colliders



### Virtual Gravitons

Fermion or VB pairs at hadron or  $e^+e^-$  colliders



# LED in monojets at D0

## Selection

- $N_{\text{jets}} = 1$  or 2
- $E_T(j_1) > 150$  GeV,  $E_T(j_2) < 50$  GeV
- $j_1$  central (good quality)
- $E_T^{\text{miss}} > 150$  GeV
- Reject events with isol. muons,
- Reject cosmics

## Results


WZ background =  $30.2 \pm 4.0$

QCD background =  $7.9 \pm 7.1$

Expected Events =  $38.0 \pm 8.2$

Observed Events = 38


$$\int L dt = 78.8 \text{ pb}^{-1}$$



# LED in monojets at D0

Limits

95% CL Limits



$n=4$

$M_D > 698$  (840) GeV

D0 DELPHI

Conservative Limit  
(no K-factor)

$n=6$

$M_D > 632$  (580) GeV

D0 DELPHI



# LED in diphotons at CDF

## Signal

$$q\bar{q} \rightarrow G_{KK} \rightarrow \gamma\gamma$$

## Backgrounds

$\gamma$ +jets


QCD dijets

## Selection

- Isolated photons
- $E_T(\gamma) > 22 \text{ GeV}$ ,  $|\eta(\gamma_1)| < 1.0$

## Results

	Central	EndCap
Observed	287	192
Expected	$280 \pm 70$	$208 \pm 42$
SM diphotons	$96 \pm 31$	$76 \pm 31$
Bkgd from fakes	$184 \pm 63$	$132 \pm 28$




# LED in diphotons at CDF

- Cross section of:

$$p\bar{p} \rightarrow q\bar{q} / gg \rightarrow \gamma\gamma + X$$

$$\frac{d\sigma}{dM_{\gamma\gamma}} = \left. \frac{d\sigma}{dM_{\gamma\gamma}} \right|_{SM} + \eta \left. \frac{d\sigma}{dM_{\gamma\gamma}} \right|_{INT} + \eta^2 \left. \frac{d\sigma}{dM_{\gamma\gamma}} \right|_{KK}$$

CDF Preliminary



- Free parameter to fit:

$$\eta = \frac{\lambda}{M_D^4}$$

Limits

*CDF*

$$M_D > 1.01 TeV$$

(This analysis)

*D0*

$$M_D > 1.21 TeV$$

- Add  $\cos\theta^*$  to fit,  $127\text{pb}^{-1}$
- Ref: PRL 86 (2001)

95% CL Limits

# Conclusions

- New Analyses of Run I Data
- Better sensitivity:
  - Improved Selection Techniques
  - Use of Full Calorimeter Acceptance
- Resulting in:
  - Improved Limits
  - Complementary and Competitive wrt LEP II
- Next Moves Have Just Started in Run II:
  - Slightly Higher Energy
  - Higher Luminosity
  - Upgraded Detectors
  - Improved Trigger Designs ... News Soon

Different theoretical notations

$$\lambda_{GRW} = \frac{-2}{\pi} \lambda_{Hew.}$$

$\lambda_{Hew.} = O(\pm 1)$       Dimensionless parameter  
Sign: destructive or constructive interference

# Prospects for Run II

## mSUGRA search in dileptons at CDF

- Estimated limit:  $m_{\tilde{g}} > 181 - 182 \text{ GeV} \rightarrow m_{\tilde{g}} > 280 - 295 \text{ GeV}$   
(same observed and expected events than Run I analysis, with  $2 \text{ fb}^{-1}$ )

## LED search in monojets at D0

- $n=2$ : sensitive to  $M_D=920 \rightarrow 1400 \text{ GeV}$
- $n=5$ : sensitive to  $M_D=700 \rightarrow 900 \text{ GeV}$   
(with  $2 \text{ TeV}$  and  $300 \text{ pb}^{-1}$ )